

pharmacy technician math calculations

The Essential Guide to Pharmacy Technician Math Calculations

pharmacy technician math calculations are the bedrock of safe and accurate medication dispensing. As a vital member of the healthcare team, a pharmacy technician must possess a strong command of various mathematical principles to ensure patient safety and efficacy of treatments. From calculating dosages and drip rates to converting units and understanding ratios, these calculations are not merely academic exercises but critical skills that directly impact patient outcomes. This comprehensive guide will delve into the core mathematical concepts every pharmacy technician needs to master, covering everything from basic arithmetic to more complex pharmaceutical calculations. We'll explore the different types of calculations, provide practical examples, and highlight why precision is paramount in this field. Mastering these skills will not only enhance your confidence in the pharmacy setting but also solidify your role as an indispensable asset to pharmacists and patients alike. Let's embark on this essential journey to understanding pharmacy math.

Table of Contents

Understanding Basic Arithmetic and Fractions

Dosage Calculations: The Cornerstone of Pharmacy Math

Understanding Ratios and Proportions

Apothecaries' and Metric Systems: Conversions You Need to Know

Understanding Percentages and Solutions

Calculating Intravenous (IV) Flow Rates

Reconstitution Calculations

Compounding Calculations: Precision in Practice

Understanding Young's Rule and Clark's Rule for Pediatric Dosing

Common Pitfalls and How to Avoid Them

Understanding Basic Arithmetic and Fractions

Before we dive into the specialized calculations specific to pharmacy, it's crucial to have a firm grasp of fundamental arithmetic. This includes proficiency with whole numbers, decimals, and, perhaps most importantly, fractions. Fractions are ubiquitous in pharmacy, representing parts of a whole, such as a fraction of a milligram or a fraction of a liter. Understanding how to add, subtract, multiply, and divide fractions, including improper fractions and mixed numbers, is non-negotiable. For instance, if a prescription calls for $\frac{1}{2}$ teaspoon and another for $\frac{1}{4}$ teaspoon, you need to be able to add these quantities correctly to determine the total volume to administer. This basic numeracy forms the foundation upon which all other pharmacy calculations are built.

Fractions often appear in prescription orders, especially when dealing with liquid medications or when specifying a portion of a larger unit. Think about situations where you might have to add or subtract fractional amounts of medication to reach a target dose. For example, if a child needs $\frac{3}{4}$ of a tablet and you only have $\frac{1}{2}$ tablet remaining, you'll need to figure out how much more is needed. Similarly, when interpreting units of measurement, understanding the relationships between different fractional parts of a whole is key. This isn't just about getting the right number; it's about ensuring the correct amount of medication is dispensed, safeguarding patient health.

Dosage Calculations: The Cornerstone of Pharmacy Math

Dosage calculations are arguably the most critical aspect of pharmacy technician mathematics. Mistakes here can have serious consequences, making accuracy paramount. These calculations typically involve determining the correct amount of medication to administer based on patient weight, age, or the concentration of the drug. You'll frequently encounter formulas that help standardize these calculations, ensuring consistency and safety across different prescriptions and patients.

One of the most common dosage calculation methods involves using the formula: $\text{Desired Dose} / \text{Have on Hand} \times \text{Volume} = \text{Amount to Administer}$. This formula is incredibly versatile. For example, if a doctor orders 250 mg of a medication, and the available stock is 500 mg per 5 mL, you would set up

the calculation as $250 \text{ mg} / 500 \text{ mg} \times 5 \text{ mL}$. Solving this reveals that you need to administer 2.5 mL of the medication. This methodical approach minimizes errors and ensures that the patient receives the precise milligram amount ordered.

Another frequent scenario involves calculating doses based on body weight, especially for pediatric or certain adult medications. The order might specify a dose per kilogram of body weight (e.g., 5 mg/kg). If a patient weighs 60 kg, the total dose would be $60 \text{ kg} \times 5 \text{ mg/kg} = 300 \text{ mg}$. You then use the "Have on Hand" formula to determine the volume of the liquid medication or the number of tablets/capsules needed to deliver that 300 mg. Understanding how to convert pounds to kilograms is also a vital skill in this context, as many weight-based orders are given in metric units.

Weight-Based Dosing

Weight-based dosing is a standard practice for many medications, ensuring that the therapeutic effect is achieved without causing toxicity. This is particularly important for children, whose metabolic rates and drug clearance can vary significantly with weight. The general principle is to calculate the total daily dose based on the prescribed milligram per kilogram (mg/kg) ratio, and then divide that dose into appropriate administration intervals (e.g., every 8 hours).

For example, a pediatrician might order Amoxicillin 20 mg/kg/day for a child weighing 44 pounds. First, you'd convert pounds to kilograms: $44 \text{ lb} / 2.2 \text{ lb/kg} = 20 \text{ kg}$. Then, calculate the total daily dose: $20 \text{ kg} \times 20 \text{ mg/kg} = 400 \text{ mg/day}$. If the Amoxicillin is supplied as 125 mg/5 mL, and the order is to administer this dose every 8 hours, you'd first determine the amount per dose: $400 \text{ mg/day} \div 3 \text{ doses/day} = 133.3 \text{ mg per dose}$. Finally, calculate the volume: $133.3 \text{ mg} / 125 \text{ mg} \times 5 \text{ mL} = \text{approximately } 5.3 \text{ mL per dose}$. This multi-step process requires careful attention to detail.

Body Surface Area (BSA) Dosing

In some specialized areas, such as oncology, medications are dosed based on Body Surface Area (BSA), a measure that accounts for both weight and height. This method is often used for potent

chemotherapy drugs where precise dosing is critical for efficacy and minimizing side effects. While the pharmacy technician may not always perform the initial BSA calculation (often done by nurses or pharmacists using nomograms or formulas), they will certainly encounter prescriptions that require calculations based on a BSA value.

The calculation typically looks like: $\text{Drug Concentration} \times \text{BSA} = \text{Dose}$. For instance, if a chemotherapy drug is ordered at 100 mg/m^2 and the patient's BSA is 1.8 m^2 , the dose would be $100 \text{ mg/m}^2 \times 1.8 \text{ m}^2 = 180 \text{ mg}$. The technician would then need to determine the volume of the drug product to prepare or dispense based on the available concentration. This type of calculation highlights the need for understanding complex drug regimens and specialized dosing protocols within pharmacy practice.

Understanding Ratios and Proportions

Ratios and proportions are fundamental mathematical concepts that are heavily utilized in pharmacy calculations, particularly in areas like dilutions, concentrations, and preparing solutions. A ratio compares two quantities, while a proportion states that two ratios are equal. Mastering these allows you to solve for unknown quantities when a relationship between them is known.

For example, if you have a concentrated stock solution and need to dilute it to a specific strength, you'll use ratios and proportions. Let's say you have a 1:10 stock solution of a disinfectant and need to prepare 500 mL of a 1:50 solution. You can set up a proportion: $(1 \text{ part solute} / 10 \text{ parts total solution}) = (x \text{ parts solute} / 500 \text{ parts total solution})$. Solving for x would tell you the amount of the stock solution needed. This concept extends to numerous pharmaceutical applications where precise dilutions are essential for drug stability and therapeutic effect.

Solving for Unknowns

The power of proportions lies in their ability to solve for an unknown variable. The classic cross-multiplication method is your best friend here. If you have a proportion like $a/b = c/d$, you can solve for any missing variable by cross-multiplying: $ad = bc$. In a pharmacy context, this could be applying it to

determine the amount of active ingredient in a given volume of a liquid medication. If a suspension contains 250 mg of active ingredient in every 5 mL, and you need to dispense 10 mL, you can set up the proportion: $250 \text{ mg} / 5 \text{ mL} = x \text{ mg} / 10 \text{ mL}$. Cross-multiplying gives you $250 \text{ mg} \times 10 \text{ mL} = 5 \text{ mL} \times x \text{ mg}$, leading to $2500 = 5x$, and therefore $x = 500 \text{ mg}$. This demonstrates how proportions can directly translate into dispensing the correct amount of medication.

Apothecaries' and Metric Systems: Conversions You Need to Know

Pharmacy historically relied heavily on the apothecaries' system, and while the metric system is now the standard in most healthcare settings, understanding apothecaries' units is still important for interpreting older prescriptions or specific formulations. Familiarity with conversions between these systems and within them (e.g., grains to grams, minims to milliliters, ounces to milliliters) is absolutely crucial for accurate dispensing. You'll need to know key equivalents like 1 grain $\approx$ 64.8 mg, 1 fluid ounce $\approx$ 30 mL, and 1 minim $\approx$ 0.06 mL.

The metric system, based on powers of 10, is more straightforward. Units like milligrams (mg), grams (g), micrograms (mcg), milliliters (mL), and liters (L) are commonly used. The ability to move between these units seamlessly is vital. For example, if a prescription is written in grams but the medication is available in milligrams, you must be able to convert 2 grams to 2000 milligrams accurately. This often involves multiplying or dividing by 1000, depending on the direction of the conversion. Precision in these conversions prevents under- or overdosing.

Metric to Apothecaries' Conversion

While the metric system is dominant, you will occasionally encounter prescriptions or dosage forms that still use apothecaries' units, especially in compounding or when referencing older drug literature. For instance, a prescriber might order 1/4 grain of a medication. As a pharmacy technician, you'll need to convert this to milligrams to accurately measure it out, especially if the available dosage forms are

in metric units. Knowing that 1 grain is approximately 64.8 mg means 1/4 grain is about 16.2 mg. This conversion is vital for ensuring the patient receives the intended dose, bridging the gap between older and modern measurement systems.

Apothecaries' to Metric Conversion

Conversely, you might need to convert apothecaries' measurements to metric units. If a prescription calls for 2 fluid ounces of a liquid, and the standard conversion is 1 fluid ounce = 30 mL, then 2 fluid ounces would be $2 \times 30 \text{ mL} = 60 \text{ mL}$. This type of conversion is common when dealing with liquid preparations or when transferring quantities between different measurement frameworks.

Understanding these equivalencies allows you to accurately prepare or dispense medications regardless of the system used in the original order.

Understanding Percentages and Solutions

Percentages are frequently used in pharmacy to express concentrations of solutions and mixtures. A percentage represents a fraction out of 100. In pharmacy, "percent by weight" (% w/w), "percent by volume" (% v/v), and "percent weight by volume" (% w/v) are all important concepts. For example, a 5% dextrose solution (% w/v) means there are 5 grams of dextrose in every 100 mL of solution.

Understanding these different types of percentages is crucial for accurate calculations. A % w/w solution indicates grams of solute per 100 grams of solution. A % v/v solution indicates milliliters of solute per 100 milliliters of solution. A % w/v solution indicates grams of solute per 100 milliliters of solution. Each has specific implications for how you calculate the amount of drug needed or the volume of a preparation to dispense.

Weight/Weight (w/w) Calculations

Weight/weight percentage calculations are most common in compounding solid dosage forms or

ointments where the density of components might vary. A 10% w/w ointment, for example, means that for every 100 grams of ointment, 10 grams are the active drug, and 90 grams are the ointment base. If you need to prepare 50 grams of this ointment, you would calculate: 10% of 50 grams = $0.10 \times 50 \text{ g} = 5 \text{ grams}$ of active drug. The remaining 45 grams would be the ointment base. Precision here ensures the correct concentration of the active ingredient in the final product.

Weight/Volume (w/v) Calculations

Weight/volume percentage is very common for liquid solutions, such as intravenous fluids or oral suspensions. A 2% w/v solution means 2 grams of solute are dissolved in enough solvent to make 100 mL of solution. If a doctor orders 1000 mL of a 0.9% w/v sodium chloride solution (normal saline), you know this concentration translates to 0.9 grams of sodium chloride per 100 mL of solution. To prepare 1000 mL, you would set up a proportion: $(0.9 \text{ g} / 100 \text{ mL}) = (x \text{ g} / 1000 \text{ mL})$. Solving this gives you $x = 9 \text{ grams}$ of sodium chloride needed. This is a fundamental calculation for preparing sterile intravenous admixtures.

Volume/Volume (v/v) Calculations

Volume/volume percentage is typically used when both the solute and the solvent are liquids. A 5% v/v solution means there are 5 mL of solute for every 100 mL of total solution. If you need to prepare 250 mL of a 5% v/v solution of alcohol, you would calculate the volume of alcohol needed: 5% of 250 mL = $0.05 \times 250 \text{ mL} = 12.5 \text{ mL}$ of alcohol. The remaining volume, $250 \text{ mL} - 12.5 \text{ mL} = 237.5 \text{ mL}$, would be the diluent (e.g., water). This calculation is essential for preparing solutions where the volume of the active ingredient contributes to the final volume.

Calculating Intravenous (IV) Flow Rates

Calculating IV flow rates is a critical skill for pharmacy technicians involved in sterile compounding and admixture services. This involves determining how quickly an IV solution should be infused over a

specific period. The primary formula used is: Total Volume to Infuse / Time to Infuse = Flow Rate. However, the units of time can vary (hours or minutes), and the IV administration set often has a calibration in drops per milliliter (gtts/mL), necessitating further calculations.

If an IV bag contains 1000 mL of fluid to be infused over 8 hours, the flow rate would be $1000 \text{ mL} / 8 \text{ hours} = 125 \text{ mL/hour}$. This is a straightforward volume per time calculation. However, if you need to determine the drip rate in drops per minute, you'll need to use the drop factor of the tubing. For example, if the tubing has a drop factor of 20 gtts/mL, you would calculate the drip rate: $(\text{Flow Rate in mL/hr} \times \text{Drop Factor}) / 60 \text{ minutes/hour} = \text{Drip Rate in gtts/min}$. So, $(125 \text{ mL/hr} \times 20 \text{ gtts/mL}) / 60 \text{ min/hr} = 2500 \text{ gtts/min} / 60 \text{ min/hr} = \text{approximately } 41.7 \text{ drops per minute}$.

Using Drop Factors

The drop factor of an IV administration set is crucial when calculating drip rates for gravity infusions. These sets come with different calibrations, commonly 10, 15, 20, or 60 drops per milliliter (gtts/mL). A higher drop factor means fewer drops are needed to deliver a milliliter of fluid, leading to a faster drip rate for the same volume. For instance, a 60 gtts/mL set is often used for very small volumes or for pediatric infusions where precise, small aliquots are administered.

The formula for calculating drip rate using the drop factor is: $(\text{Total Volume} \times \text{Drop Factor}) / \text{Total Time in Minutes} = \text{Drip Rate (gtts/min)}$. Let's say you need to infuse 500 mL of fluid over 4 hours using a 15 gtts/mL administration set. First, convert 4 hours to minutes: $4 \text{ hours} \times 60 \text{ minutes/hour} = 240 \text{ minutes}$. Then, plug into the formula: $(500 \text{ mL} \times 15 \text{ gtts/mL}) / 240 \text{ min} = 7500 \text{ gtts} / 240 \text{ min} = 31.25 \text{ gtts/min}$. This tells you the IV should be set to drip at approximately 31 drops per minute.

Reconstitution Calculations

Many antibiotics and other medications are supplied as powders that must be reconstituted with a specific diluent (like sterile water or saline) before they can be administered. Reconstitution calculations involve determining the correct amount of diluent to add and then calculating the final

concentration of the reconstituted drug. This ensures the medication is in a stable, administrable form at the correct strength.

The manufacturer's instructions are your primary guide. They will specify the volume of diluent to add and the resulting concentration. For example, a vial of cefazolin powder might state: "Add 10 mL sterile water for injection for a final concentration of approximately 225 mg/mL." Once reconstituted, if a prescription calls for 500 mg of cefazolin, you would calculate the volume to draw up: $(\text{Desired Dose} / \text{Concentration}) = \text{Volume to Administer}$. So, $(500 \text{ mg} / 225 \text{ mg/mL}) =$ approximately 2.2 mL. It is crucial to check the stability and storage requirements after reconstitution, as these medications often have limited shelf lives.

Determining Final Concentration

Before calculating the volume to administer, you must first establish the final concentration after reconstitution. Manufacturers provide specific instructions. For instance, if a vial of vancomycin powder states "Reconstitute with 20 mL sterile water for injection to yield approximately 50 mg/mL," then that is your final concentration. If a prescription is for 750 mg of vancomycin, the calculation is straightforward: $(\text{Desired Dose} / \text{Concentration}) = \text{Volume to draw}$. Therefore, $(750 \text{ mg} / 50 \text{ mg/mL}) =$ 15 mL. Always read and follow the instructions on the drug's packaging meticulously.

Compounding Calculations: Precision in Practice

Compounding is the art and science of preparing customized medications tailored to an individual patient's needs. This can involve mixing multiple ingredients, adjusting dosages, or preparing non-sterile or sterile formulations. Pharmacy technicians play a crucial role in compounding, and accurate calculations are paramount to ensure the safety, efficacy, and quality of the final product. This area demands meticulous attention to detail and a solid understanding of ratios, percentages, and unit conversions.

When compounding, you might need to calculate the amount of each active ingredient and excipient

(inactive ingredient) required for a specific batch size. This often involves using proportions, alligation, or percentage calculations, depending on the nature of the ingredients and the desired final product. For example, if you're compounding an ointment with a specific percentage of an active drug, you'll need to calculate the exact weight of that drug and the ointment base required for the total weight of the ointment.

Alligation Medial and Alternate

Alligation is a mathematical method used in pharmacy for calculating the amounts of different strengths of a drug or solution needed to obtain a desired intermediate strength. There are two main types: alligation medial and alligation alternate. Alligation alternate is more commonly used by technicians. It helps determine the ratio of two strengths of a drug needed to create a mixture of a specific concentration.

Here's a simplified explanation of alligation alternate: You want to mix a higher concentration drug with a lower concentration drug to achieve a desired intermediate concentration. You set up a grid with the desired strength in the center. On one diagonal, you subtract the strengths (higher minus lower). The resulting numbers represent the "parts" of each ingredient you need. For example, to mix a 10% solution and a 30% solution to get a 20% solution: Higher strength (30%) - Desired strength (20%) = 10 parts of the 10% solution. Desired strength (20%) - Lower strength (10%) = 10 parts of the 30% solution. So, you'd mix equal parts of the 10% and 30% solutions to get a 20% solution. This method is invaluable for accurate compounding.

Understanding Young's Rule and Clark's Rule for Pediatric Dosing

Dosage calculations for children are particularly important because their bodies metabolize and excrete drugs differently than adults. Two common methods used to calculate pediatric dosages based on adult dosages are Young's Rule and Clark's Rule. While the pharmacist is typically responsible for

determining the appropriate dosing method, pharmacy technicians may encounter prescriptions that require these calculations or need to verify them.

Young's Rule is based on a child's age in years. The formula is: $(\text{Child's Age in Years} / \text{Adult Dose}) \times \text{Adult Dose} = \text{Pediatric Dose}$. For example, if an adult dose of a medication is 500 mg and the child is 5 years old, the pediatric dose would be $(5 / 20) \times 500 \text{ mg} = 125 \text{ mg}$. It's important to note that Young's Rule is generally used for children over the age of two.

Clark's Rule

Clark's Rule is another method for calculating pediatric doses, and it is based on the child's weight. This is often preferred as weight is a more reliable indicator of drug dosage than age, especially for infants and younger children. The formula is: $(\text{Child's Weight in Pounds} / 150 \text{ lbs}) \times \text{Adult Dose} = \text{Pediatric Dose}$. For instance, if the adult dose is 300 mg and the child weighs 50 pounds, the pediatric dose would be $(50 \text{ lbs} / 150 \text{ lbs}) \times 300 \text{ mg} = 100 \text{ mg}$. Always remember to convert pounds to kilograms if the weight is given in metric units, using the factor $1 \text{ kg} = 2.2 \text{ lbs}$, when using formulas that require weight in pounds.

Common Pitfalls and How to Avoid Them

Even with strong foundational math skills, errors can occur in pharmacy calculations. Recognizing common pitfalls is the first step toward preventing them. One of the most frequent mistakes is simple transposing numbers or decimal point errors, which can lead to a tenfold or even a hundredfold error in dosage. Double-checking your work is not just a good practice; it's a necessity.

Another pitfall is not paying attention to the units of measurement. If an order is in milligrams but your medication is in grams, or if you're mixing metric and apothecaries' units without proper conversion, errors are almost guaranteed. Always verify that the units in your calculation match the units of the medication you are preparing or dispensing. Taking the time to write down each step of your calculation clearly can help you catch these errors before they impact patient care. Furthermore, never

guess; if you are unsure about a calculation, always ask a pharmacist for verification.

Unit Errors

Unit errors are incredibly common and can lead to significant dosing mistakes. For example, confusing milligrams (mg) with grams (g) or milliliters (mL) with liters (L) can result in a dose that is 1000 times too large or too small. Always write out the units for each number in your calculation and ensure they cancel out correctly. If you are converting between units, be sure you are using the correct conversion factor. For example, converting 1000 mcg to milligrams requires dividing by 1000 ($1000 \text{ mcg} / 1000 \text{ mcg/mg} = 1 \text{ mg}$), not multiplying.

Decimal Point Errors

Decimal point errors are perhaps the most insidious. A misplaced decimal can turn a safe dose into a toxic one, or a therapeutic dose into an ineffective one. When calculating, carefully align your decimal points. If you are working with a calculator, take an extra moment to confirm the number you've entered. For manual calculations, it can be helpful to estimate your answer beforehand. For example, if you are calculating a dose and your estimate is around 5 mL, but your calculation yields 50 mL or 0.5 mL, that's a strong indicator that you need to recheck your work for a decimal point error.

These calculations are not just about numbers; they are about patient safety. The skills you hone in mastering pharmacy math calculations directly translate into better patient care. By understanding the principles, practicing diligently, and always prioritizing accuracy, you can confidently perform these essential tasks. Remember, a pharmacy technician's role is indispensable, and a strong grasp of mathematics is a key component of that crucial role. Keep practicing, keep learning, and always, always double-check your work. Your commitment to accuracy makes a real difference in people's lives every single day.

FAQ

Q: What are the most common types of math calculations a pharmacy technician performs?

A: Pharmacy technicians commonly perform dosage calculations (using formulas like $\text{Desired/Have} \times \text{Quantity}$), unit conversions (metric, apothecaries', household), ratio and proportion calculations for dilutions and concentrations, IV flow rate calculations, reconstitution calculations for powdered medications, and percentage calculations for solutions and ointments.

Q: Why is it important to understand both the metric and apothecaries' systems?

A: While the metric system is the standard in modern pharmacy, older prescriptions, certain drug references, and compounding practices may still utilize apothecaries' units. Understanding both allows technicians to accurately interpret all types of medication orders and ensure the correct dosage is dispensed, preventing potentially dangerous errors.

Q: How do you calculate pediatric doses when an adult dose is known?

A: Pediatric doses are often calculated using methods like Young's Rule (based on age) or Clark's Rule (based on weight). Clark's Rule, which uses the child's weight in pounds divided by 150 lbs, multiplied by the adult dose, is generally considered more accurate as weight is a better indicator of drug metabolism than age.

Q: What is the importance of checking the "have on hand" amount

when calculating drug dosages?

A: The "have on hand" amount represents the concentration or form of the medication that is available in the pharmacy (e.g., 250 mg per 5 mL of a liquid, or 10 mg per tablet). This value is critical for determining the volume or number of units (mL, tablets) the patient needs to receive to obtain the desired dose. Without it, you cannot accurately calculate the final administration quantity.

Q: What is reconstitution, and what kind of calculations are involved?

A: Reconstitution is the process of adding a diluent (like sterile water or saline) to a powdered medication to dissolve it and make it administrable, typically as a liquid. Calculations involve determining the correct volume of diluent to add according to the manufacturer's instructions to achieve a specific final concentration (e.g., mg/mL), and then calculating the volume of this reconstituted solution that contains the prescribed dose.

Q: How are percentages used in pharmacy math calculations?

A: Percentages are used to express concentrations of solutions, ointments, and other preparations. Common types include weight/weight (% w/w), volume/volume (% v/v), and weight/volume (% w/v). For example, a 5% w/v solution means 5 grams of solute in 100 mL of solution. Technicians use these to calculate the amount of active ingredient needed for a specific preparation.

Q: What is an IV flow rate calculation, and why is it important?

A: IV flow rate calculations determine how quickly an intravenous solution should be administered. They involve calculating the volume per unit of time (e.g., mL/hour) and, when using gravity drip sets, the number of drops per minute (using the set's drop factor). Accurate flow rates are essential for ensuring the medication is delivered at the prescribed rate to achieve therapeutic effect and avoid complications like under-infusion or fluid overload.

Q: What are some common mistakes pharmacy technicians make in math calculations, and how can they be avoided?

A: Common mistakes include unit errors (confusing mg with g, mL with L), decimal point errors, and misinterpreting the "have on hand" quantity. To avoid these, technicians should always double-check their work, write out units clearly, perform estimations before detailed calculations, and never hesitate to ask a pharmacist for verification if unsure.

[Pharmacy Technician Math Calculations](#)

Pharmacy Technician Math Calculations

Related Articles

- [pure math jobs](#)
- [radical math examples](#)
- [represent definition math](#)

[Back to Home](#)