

WHAT IS FREE FALL MOTION IN PHYSICS

UNDERSTANDING WHAT IS FREE FALL MOTION IN PHYSICS

WHAT IS FREE FALL MOTION IN PHYSICS IS A FUNDAMENTAL CONCEPT THAT DESCRIBES THE MOVEMENT OF AN OBJECT SOLELY UNDER THE INFLUENCE OF GRAVITY. IT'S A CAPTIVATING PHENOMENON THAT EXPLAINS WHY OBJECTS FALL TOWARDS THE EARTH, FROM A DROPPED APPLE TO THE ORBITS OF CELESTIAL BODIES. THIS ARTICLE WILL DELVE DEEP INTO THE NUANCES OF FREE FALL, EXPLORING ITS DEFINITION, THE FORCES INVOLVED, THE EFFECTS OF GRAVITY, AND THE PHYSICS PRINCIPLES THAT GOVERN THIS SEEMINGLY SIMPLE YET COMPLEX MOTION. WE WILL UNRAVEL THE EQUATIONS THAT DESCRIBE FREE FALL, DISTINGUISH IT FROM PROJECTILE MOTION, AND EVEN TOUCH UPON ITS PRESENCE IN REAL-WORLD SCENARIOS AND SPACE EXPLORATION.

TABLE OF CONTENTS

WHAT IS FREE FALL MOTION IN PHYSICS?

THE DEFINING CHARACTERISTICS OF FREE FALL

GRAVITY: THE UNSEEN FORCE

EQUATIONS GOVERNING FREE FALL MOTION

FREE FALL VS. PROJECTILE MOTION: A CRUCIAL DISTINCTION

REAL-WORLD EXAMPLES OF FREE FALL

FREE FALL IN SPACE AND BEYOND

COMMON MISCONCEPTIONS ABOUT FREE FALL

WHAT IS FREE FALL MOTION IN PHYSICS?

AT ITS CORE, FREE FALL MOTION IN PHYSICS REFERS TO THE MOTION OF AN OBJECT WHERE THE ONLY FORCE ACTING UPON IT IS GRAVITY. IMAGINE DROPPING A BALL; AS SOON AS IT LEAVES YOUR HAND, AND WE IGNORE AIR RESISTANCE FOR A MOMENT, GRAVITY IS THE SOLE PROPRIETOR OF ITS DOWNWARD JOURNEY. THIS IDEALIZED SCENARIO SIMPLIFIES THE COMPLEX INTERACTIONS THAT OFTEN OCCUR IN THE PHYSICAL WORLD, ALLOWING US TO ISOLATE AND UNDERSTAND THE FUNDAMENTAL PRINCIPLES AT PLAY. IT'S A CORNERSTONE CONCEPT IN CLASSICAL MECHANICS, ESSENTIAL FOR COMPREHENDING EVERYTHING FROM THE TRAJECTORY OF A THROWN OBJECT TO THE VAST CELESTIAL BALLET OF PLANETS AND STARS.

UNDERSTANDING FREE FALL IS CRUCIAL BECAUSE IT FORMS THE BASIS FOR MANY OTHER PHYSICS PRINCIPLES. WITHOUT GRASPING HOW GRAVITY AFFECTS OBJECTS IN ISOLATION, IT BECOMES CHALLENGING TO ANALYZE MORE COMPLEX SYSTEMS. WE OFTEN SIMPLIFY REAL-WORLD SITUATIONS BY ASSUMING FREE FALL CONDITIONS TO MAKE CALCULATIONS AND PREDICTIONS MANAGEABLE. THIS ALLOWS US TO BUILD A FOUNDATIONAL UNDERSTANDING THAT CAN THEN BE EXPANDED UPON TO ACCOUNT FOR ADDITIONAL FORCES AND COMPLEXITIES.

THE DEFINING CHARACTERISTICS OF FREE FALL

SEVERAL KEY CHARACTERISTICS DEFINE FREE FALL MOTION. THE MOST CRITICAL IS THE ABSENCE OF ANY FORCES OTHER THAN GRAVITY. THIS MEANS WE ARE NEGLECTING EXTERNAL INFLUENCES LIKE AIR RESISTANCE, THRUST, OR FRICTION. IN A VACUUM, ANY OBJECT, REGARDLESS OF ITS MASS OR COMPOSITION, WOULD EXPERIENCE THE SAME ACCELERATION DUE TO GRAVITY. THIS IS A PROFOUND INSIGHT, FAMOUSLY DEMONSTRATED BY GALILEO GALILEI, WHO THEORIZED THAT OBJECTS OF DIFFERENT WEIGHTS FALL AT THE SAME RATE. THE ONLY FORCE PULLING THEM DOWN IS THE SAME GRAVITATIONAL PULL, CAUSING THEM TO ACCELERATE EQUALLY.

ANOTHER DEFINING CHARACTERISTIC IS THE CONSTANT ACCELERATION EXPERIENCED BY THE OBJECT. ON EARTH'S SURFACE, THIS ACCELERATION DUE TO GRAVITY IS APPROXIMATELY 9.8 METERS PER SECOND SQUARED (m/s^2), OFTEN DENOTED BY THE SYMBOL 'G'. THIS MEANS THAT FOR EVERY SECOND AN OBJECT IS IN FREE FALL, ITS DOWNWARD VELOCITY INCREASES BY 9.8 M/S. THIS CONSTANT RATE OF CHANGE IN VELOCITY IS WHAT MAKES FREE FALL MOTION PREDICTABLE AND CALCULABLE USING KINEMATIC EQUATIONS. THIS CONSISTENT ACCELERATION IS A DIRECT CONSEQUENCE OF THE GRAVITATIONAL FORCE BEING

RELATIVELY UNIFORM OVER THE SMALL DISTANCES INVOLVED IN MOST EVERYDAY FREE FALL SCENARIOS.

THE ROLE OF MASS IN FREE FALL

A COMMON MISCONCEPTION IS THAT HEAVIER OBJECTS FALL FASTER THAN LIGHTER ONES. HOWEVER, IN A TRUE FREE FALL SCENARIO, MASS PLAYS NO ROLE IN THE ACCELERATION. GALILEO'S EXPERIMENTS, AND LATER CONFIRMED BY ASTRONAUTS ON THE MOON DROPPING A FEATHER AND A HAMMER SIMULTANEOUSLY, ILLUSTRATE THIS POINT. WHILE THE FORCE OF GRAVITY MIGHT BE STRONGER ON A MORE MASSIVE OBJECT (AS FORCE EQUALS MASS TIMES ACCELERATION, $F=ma$), THE INCREASED MASS ALSO MEANS IT REQUIRES MORE FORCE TO ACCELERATE THAT OBJECT AT THE SAME RATE. THESE TWO EFFECTS CANCEL EACH OTHER OUT, RESULTING IN THE SAME ACCELERATION FOR ALL OBJECTS IN FREE FALL, IRRESPECTIVE OF THEIR MASS. THIS EQUALITY OF ACCELERATION IS A FUNDAMENTAL ASPECT OF GRAVITY AND INERTIAL MASS.

INITIAL VELOCITY IN FREE FALL

THE CONCEPT OF INITIAL VELOCITY IS ALSO IMPORTANT WHEN DISCUSSING FREE FALL. AN OBJECT CAN BE IN FREE FALL WHETHER IT IS DROPPED FROM REST, THROWN DOWNWARDS, OR EVEN THROWN UPWARDS. IF AN OBJECT IS DROPPED FROM REST, ITS INITIAL VERTICAL VELOCITY IS ZERO. IF IT'S THROWN DOWNWARDS, ITS INITIAL VERTICAL VELOCITY IS POSITIVE (ASSUMING DOWNWARD IS POSITIVE). IF IT'S THROWN UPWARDS, ITS INITIAL VERTICAL VELOCITY IS NEGATIVE (IF DOWNWARD IS POSITIVE). IN ALL THESE CASES, ONCE THE OBJECT IS NO LONGER IN CONTACT WITH THE SOURCE OF ITS INITIAL MOTION (LIKE YOUR HAND) AND IS ONLY UNDER THE INFLUENCE OF GRAVITY, IT IS CONSIDERED TO BE IN FREE FALL. THE TRAJECTORY WILL DIFFER BASED ON THE INITIAL VELOCITY, BUT THE ACCELERATION DUE TO GRAVITY REMAINS CONSTANT.

GRAVITY: THE UNSEEN FORCE

GRAVITY IS THE FUNDAMENTAL FORCE THAT DICTATES FREE FALL MOTION. IT IS A UNIVERSAL FORCE OF ATTRACTION THAT EXISTS BETWEEN ANY TWO OBJECTS WITH MASS. THE MORE MASSIVE AN OBJECT, THE STRONGER ITS GRAVITATIONAL PULL. THE CLOSER TWO OBJECTS ARE, THE STRONGER THE GRAVITATIONAL ATTRACTION BETWEEN THEM. THIS IS WHY WE ARE PULLED TOWARDS THE EARTH; OUR PLANET HAS A SIGNIFICANT MASS, EXERTING A SUBSTANTIAL GRAVITATIONAL FORCE ON EVERYTHING ON AND NEAR ITS SURFACE. THIS FORCE IS WHAT CAUSES OBJECTS TO ACCELERATE DOWNWARDS WHEN RELEASED.

THE STRENGTH OF GRAVITY IS NOT UNIFORM ACROSS THE ENTIRE EARTH. WHILE WE OFTEN USE THE APPROXIMATE VALUE OF 9.8 m/s^2 , THE ACTUAL ACCELERATION DUE TO GRAVITY CAN VARY SLIGHTLY DEPENDING ON FACTORS LIKE ALTITUDE, LATITUDE, AND LOCAL GEOLOGICAL DENSITY. FOR INSTANCE, GRAVITY IS SLIGHTLY WEAKER AT THE TOP OF A MOUNTAIN THAN AT SEA LEVEL. HOWEVER, FOR MOST INTRODUCTORY PHYSICS PROBLEMS AND EVERYDAY CALCULATIONS, THE CONSTANT VALUE OF ' g ' IS A HIGHLY ACCURATE APPROXIMATION FOR UNDERSTANDING FREE FALL MOTION.

NEWTON'S LAW OF UNIVERSAL GRAVITATION

SIR ISAAC NEWTON'S LAW OF UNIVERSAL GRAVITATION PROVIDES A MATHEMATICAL FRAMEWORK FOR UNDERSTANDING THE FORCE OF GRAVITY. IT STATES THAT EVERY PARTICLE ATTRACTS EVERY OTHER PARTICLE IN THE UNIVERSE WITH A FORCE THAT IS DIRECTLY PROPORTIONAL TO THE PRODUCT OF THEIR MASSES AND INVERSELY PROPORTIONAL TO THE SQUARE OF THE DISTANCE BETWEEN THEIR CENTERS. MATHEMATICALLY, THIS IS EXPRESSED AS $F = G (m_1 m_2) / r^2$, WHERE F IS THE GRAVITATIONAL FORCE, G IS THE GRAVITATIONAL CONSTANT, m_1 AND m_2 ARE THE MASSES OF THE TWO OBJECTS, AND r IS THE DISTANCE BETWEEN THEIR CENTERS.

WHILE THIS LAW APPLIES UNIVERSALLY, FOR OBJECTS NEAR THE EARTH'S SURFACE, THE CALCULATION SIMPLIFIES. BECAUSE THE EARTH'S MASS (m_1) AND RADIUS (r) ARE CONSTANT FOR OBJECTS NEAR ITS SURFACE, AND THE GRAVITATIONAL CONSTANT (G) IS ALSO A FIXED VALUE, THE TERM $G m_1 / r^2$ EFFECTIVELY BECOMES OUR CONSTANT ' g '. THIS EXPLAINS WHY OBJECTS

EXPERIENCE A CONSTANT ACCELERATION DUE TO GRAVITY NEAR THE EARTH'S SURFACE. IT'S A DIRECT CONSEQUENCE OF NEWTON'S GRAND THEORY OF GRAVITY.

EQUATIONS GOVERNING FREE FALL MOTION

THE MOTION OF AN OBJECT IN FREE FALL CAN BE DESCRIBED USING A SET OF KINEMATIC EQUATIONS, WHICH ARE DERIVED FROM THE PRINCIPLES OF CONSTANT ACCELERATION. THESE EQUATIONS ALLOW US TO PREDICT AN OBJECT'S POSITION, VELOCITY, AND TIME OF FLIGHT. ASSUMING WE DEFINE THE DOWNWARD DIRECTION AS POSITIVE AND NEGLECT AIR RESISTANCE, THE PRIMARY EQUATIONS ARE:

- **$v = u + at$** : THIS EQUATION RELATES FINAL VELOCITY (v) TO INITIAL VELOCITY (u), ACCELERATION (a), AND TIME (t). IN FREE FALL, ' a ' IS REPLACED BY ' g '. SO, $v = u + gt$.
- **$s = ut + \frac{1}{2}at^2$** : THIS EQUATION DESCRIBES DISPLACEMENT (s) IN TERMS OF INITIAL VELOCITY (u), ACCELERATION (a), AND TIME (t). FOR FREE FALL, $s = ut + \frac{1}{2}gt^2$.
- **$v^2 = u^2 + 2as$** : THIS EQUATION RELATES FINAL VELOCITY (v) TO INITIAL VELOCITY (u), ACCELERATION (a), AND DISPLACEMENT (s). IN FREE FALL, $v^2 = u^2 + 2gs$.

THESE EQUATIONS ARE INCREDIBLY POWERFUL TOOLS FOR SOLVING PROBLEMS INVOLVING FREE FALL. FOR INSTANCE, IF YOU DROP AN OBJECT FROM A CERTAIN HEIGHT, YOU CAN USE THESE EQUATIONS TO CALCULATE HOW LONG IT WILL TAKE TO HIT THE GROUND OR WHAT ITS VELOCITY WILL BE UPON IMPACT. CONVERSELY, IF YOU KNOW THE TIME AN OBJECT HAS BEEN FALLING, YOU CAN CALCULATE THE DISTANCE IT HAS TRAVELED OR ITS FINAL SPEED.

CALCULATING TIME TO REACH THE GROUND

TO CALCULATE THE TIME IT TAKES FOR AN OBJECT TO REACH THE GROUND FROM A KNOWN HEIGHT, WE OFTEN USE THE EQUATION $s = ut + \frac{1}{2}gt^2$, WHERE ' s ' IS THE INITIAL HEIGHT. IF THE OBJECT IS DROPPED FROM REST, THEN ' u ' IS 0. THE EQUATION SIMPLIFIES TO $s = \frac{1}{2}gt^2$. REARRANGING THIS TO SOLVE FOR ' t ', WE GET $t = \sqrt{\frac{2s}{g}}$ ($\sqrt{2s/g}$). THIS FORMULA IS INCREDIBLY USEFUL FOR UNDERSTANDING HOW THE HEIGHT FROM WHICH AN OBJECT IS DROPPED AFFECTS ITS FALL TIME. A GREATER HEIGHT MEANS A LONGER TIME TO FALL, BUT THE RELATIONSHIP IS NOT LINEAR; IT'S DEPENDENT ON THE SQUARE ROOT OF THE HEIGHT.

DETERMINING FINAL VELOCITY

THE FINAL VELOCITY OF AN OBJECT IN FREE FALL CAN BE DETERMINED USING THE EQUATIONS $v = u + gt$ OR $v^2 = u^2 + 2gs$. IF AN OBJECT IS DROPPED FROM REST ($u=0$), THE FIRST EQUATION BECOMES $v = gt$, AND THE SECOND BECOMES $v^2 = 2gs$, OR $v = \sqrt{2gs}$ ($\sqrt{2gs}$). THIS HIGHLIGHTS HOW THE VELOCITY INCREASES LINEARLY WITH TIME, BUT WITH THE SQUARE ROOT DEPENDENCE ON THE DISTANCE FALLEN. IT'S FASCINATING TO SEE HOW THESE SIMPLE RELATIONSHIPS CAN PREDICT COMPLEX OUTCOMES.

FREE FALL VS. PROJECTILE MOTION: A CRUCIAL DISTINCTION

IT'S IMPORTANT TO DISTINGUISH BETWEEN PURE FREE FALL AND PROJECTILE MOTION, ALTHOUGH THEY ARE CLOSELY RELATED. FREE FALL, AS WE'VE DEFINED IT, IS MOTION SOLELY UNDER THE INFLUENCE OF GRAVITY. PROJECTILE MOTION, ON THE OTHER HAND, DESCRIBES THE MOTION OF AN OBJECT THAT IS LAUNCHED OR THROWN INTO THE AIR AND THEN MOVES UNDER THE INFLUENCE OF GRAVITY ALONE, AFTER THE INITIAL LAUNCH FORCE IS NO LONGER ACTING. THE KEY DIFFERENCE LIES IN THE INITIAL

CONDITIONS AND THE PRESENCE OF A HORIZONTAL COMPONENT OF VELOCITY.

IN PURE FREE FALL, THE OBJECT TYPICALLY HAS ZERO INITIAL HORIZONTAL VELOCITY. ITS MOTION IS ENTIRELY VERTICAL. IN PROJECTILE MOTION, THE OBJECT IS GIVEN AN INITIAL VELOCITY THAT HAS BOTH A HORIZONTAL AND A VERTICAL COMPONENT. THINK OF THROWING A BALL FORWARD. THE HORIZONTAL MOTION CONTINUES (IDEALLY) AT A CONSTANT SPEED BECAUSE THERE'S NO HORIZONTAL FORCE ACTING ON IT (NEGLECTING AIR RESISTANCE), WHILE THE VERTICAL MOTION IS PURE FREE FALL. THE COMBINATION OF THESE TWO INDEPENDENT MOTIONS RESULTS IN A PARABOLIC TRAJECTORY.

THE INDEPENDENCE OF HORIZONTAL AND VERTICAL MOTION

A FUNDAMENTAL PRINCIPLE IN UNDERSTANDING PROJECTILE MOTION IS THE INDEPENDENCE OF ITS HORIZONTAL AND VERTICAL COMPONENTS. THE HORIZONTAL VELOCITY OF A PROJECTILE REMAINS CONSTANT THROUGHOUT ITS FLIGHT (AGAIN, NEGLECTING AIR RESISTANCE), WHILE THE VERTICAL MOTION IS GOVERNED BY THE EQUATIONS OF FREE FALL. THIS MEANS THAT AN OBJECT FIRED HORIZONTALLY FROM A HEIGHT WILL HIT THE GROUND AT THE SAME TIME AS AN OBJECT SIMPLY DROPPED FROM THE SAME HEIGHT. THE HORIZONTAL SPEED ONLY AFFECTS HOW FAR THE OBJECT TRAVELS HORIZONTALLY BEFORE LANDING.

AIR RESISTANCE: THE REAL-WORLD COMPLICATION

IN THE REAL WORLD, VERY FEW OBJECTS EXPERIENCE PERFECT FREE FALL BECAUSE OF AIR RESISTANCE. AIR RESISTANCE IS A FORM OF FRICTION THAT OPPOSES THE MOTION OF AN OBJECT THROUGH THE AIR. IT DEPENDS ON FACTORS SUCH AS THE OBJECT'S SPEED, SHAPE, AND THE DENSITY OF THE AIR. FOR LIGHT, LARGE OBJECTS LIKE FEATHERS, AIR RESISTANCE CAN SIGNIFICANTLY SLOW THEIR DESCENT, MAKING THEM APPEAR TO FALL MUCH SLOWER THAN HEAVIER, DENSER OBJECTS. THIS IS WHY OUR IDEALIZED FREE FALL CALCULATIONS ARE OFTEN APPROXIMATIONS THAT WORK BEST IN A VACUUM OR FOR DENSE, COMPACT OBJECTS MOVING AT RELATIVELY LOW SPEEDS.

REAL-WORLD EXAMPLES OF FREE FALL

WHILE PERFECT FREE FALL IS AN IDEALIZATION, MANY REAL-WORLD SCENARIOS CLOSELY APPROXIMATE IT, ALLOWING US TO APPLY THE PRINCIPLES WE'VE DISCUSSED. ONE OF THE MOST COMMON EXAMPLES IS DROPPING AN OBJECT FROM A SIGNIFICANT HEIGHT. WHEN YOU DROP A STONE FROM A BRIDGE, THE INITIAL MOMENT IT LEAVES YOUR HAND IS THE BEGINNING OF ITS FREE FALL. THE ACCELERATION DUE TO GRAVITY PULLS IT DOWNWARDS, AND ITS SPEED INCREASES UNTIL IT REACHES THE GROUND OR ENCOUNTERS SIGNIFICANT AIR RESISTANCE.

ANOTHER EVERYDAY EXAMPLE, ALBEIT BRIEF, IS THE FEELING OF WEIGHTLESSNESS WHEN YOU GO OVER A SMALL BUMP IN A CAR OR ON A ROLLER COASTER. FOR THAT FLEETING MOMENT, YOUR BODY IS ESSENTIALLY IN FREE FALL, NOT BEING SUPPORTED BY THE SEAT. THIS SENSATION IS A DIRECT MANIFESTATION OF THE PRINCIPLES OF FREE FALL MOTION. SIMILARLY, BUNGEE JUMPING IS A DRAMATIC EXAMPLE WHERE THE JUMPER EXPERIENCES PERIODS OF FREE FALL BEFORE THE CORD BRINGS THEM TO A HALT. THE THRILL COMES FROM THE CONTROLLED DESCENT UNDER GRAVITY.

OBJECTS FALLING FROM BUILDINGS

CONSIDER AN OBJECT FALLING FROM THE TOP OF A SKYSCRAPER. ASSUMING AIR RESISTANCE IS NOT A DOMINANT FACTOR FOR A DENSE OBJECT OVER A CERTAIN RANGE OF HEIGHTS, ITS DESCENT CAN BE MODELED USING FREE FALL EQUATIONS. THIS IS WHY ARCHITECTS AND ENGINEERS NEED TO UNDERSTAND THE PHYSICS OF FALLING OBJECTS WHEN DESIGNING STRUCTURES, CONSIDERING FACTORS LIKE WIND LOADS AND POTENTIAL FALLING DEBRIS. THE IMPACT FORCE OF SUCH AN OBJECT IS DIRECTLY RELATED TO ITS VELOCITY UPON IMPACT, WHICH IS A PRODUCT OF FREE FALL ACCELERATION.

SKYDIVERS AND TERMINAL VELOCITY

SKYDIVERS ARE A FASCINATING CASE STUDY THAT BRIDGES THE GAP BETWEEN IDEALIZED FREE FALL AND REAL-WORLD PHYSICS. INITIALLY, A SKYDIVER ACCELERATES DOWNWARDS ACCORDING TO THE FREE FALL EQUATIONS. HOWEVER, AS THEIR SPEED INCREASES, AIR RESISTANCE ALSO INCREASES. EVENTUALLY, THE FORCE OF AIR RESISTANCE BECOMES EQUAL IN MAGNITUDE TO THE FORCE OF GRAVITY. AT THIS POINT, THE NET FORCE ON THE SKYDIVER IS ZERO, AND THEY STOP ACCELERATING, REACHING A CONSTANT SPEED KNOWN AS TERMINAL VELOCITY. THIS IS NOT PURE FREE FALL, BUT IT'S A DIRECT CONSEQUENCE OF THE INTERPLAY BETWEEN GRAVITY AND AIR RESISTANCE.

FREE FALL IN SPACE AND BEYOND

THE CONCEPT OF FREE FALL EXTENDS FAR BEYOND EARTH'S ATMOSPHERE AND IS CRUCIAL FOR UNDERSTANDING CELESTIAL MECHANICS. SATELLITES ORBITING THE EARTH ARE, IN ESSENCE, IN A STATE OF CONTINUOUS FREE FALL. THEY ARE CONSTANTLY FALLING TOWARDS THE EARTH, BUT THEIR HORIZONTAL VELOCITY IS SO GREAT THAT THEY CONTINUOUSLY "MISS" THE EARTH, RESULTING IN AN ORBIT. THIS ELEGANT DANCE OF GRAVITY AND INERTIA KEEPS THEM IN PERPETUAL MOTION AROUND OUR PLANET.

SIMILARLY, THE MOON IS IN FREE FALL AROUND THE EARTH, AND THE EARTH IS IN FREE FALL AROUND THE SUN. THE ORBITS OF PLANETS, STARS, AND GALAXIES ARE ALL GOVERNED BY THE PRINCIPLES OF GRAVITATIONAL ATTRACTION AND THE RESULTING FREE FALL. SPACECRAFT TRAVELING TO OTHER PLANETS ALSO UTILIZE GRAVITATIONAL FORCES, OFTEN EMPLOYING "GRAVITATIONAL ASSISTS" WHERE THEY USE THE GRAVITY OF A PLANET TO CHANGE THEIR SPEED AND DIRECTION, EFFECTIVELY RIDING ALONG A GRAVITATIONAL CURVE. THIS UNDERSTANDING ALLOWS US TO NAVIGATE THE VASTNESS OF SPACE.

ORBITAL MECHANICS

ORBITAL MECHANICS IS A DIRECT APPLICATION OF FREE FALL PRINCIPLES. AN OBJECT IN ORBIT IS CONSTANTLY ACCELERATING TOWARDS THE CENTRAL BODY DUE TO GRAVITY. HOWEVER, ITS TANGENTIAL VELOCITY (ITS SPEED PARALLEL TO THE ORBIT) PREVENTS IT FROM CRASHING INTO THE CENTRAL BODY. INSTEAD, IT FOLLOWS A CURVED PATH. THE SPEED AND ALTITUDE OF AN ORBIT DETERMINE ITS PERIOD – HOW LONG IT TAKES TO COMPLETE ONE REVOLUTION. THIS DELICATE BALANCE IS WHAT KEEPS EVERYTHING FROM THE INTERNATIONAL SPACE STATION TO DISTANT EXOPLANETS IN THEIR CELESTIAL PATHS.

WEIGHTLESSNESS AND FREE FALL

THE SENSATION OF WEIGHTLESSNESS EXPERIENCED BY ASTRONAUTS IN ORBIT IS A RESULT OF BEING IN A CONTINUOUS STATE OF FREE FALL. BOTH THE ASTRONAUTS AND THEIR SPACECRAFT ARE FALLING AROUND THE EARTH AT THE SAME RATE. BECAUSE THEY ARE FALLING TOGETHER, THE ASTRONAUTS DON'T FEEL THE PULL OF GRAVITY IN THE SAME WAY THEY WOULD ON EARTH'S SURFACE. THEY ARE ESSENTIALLY "FLOATING" WITHIN THEIR ENVIRONMENT. THIS IS A PROFOUND ILLUSTRATION THAT EVEN WHEN UNDER THE INFLUENCE OF STRONG GRAVITY, BEING IN FREE FALL CAN LEAD TO THE PERCEPTION OF NO WEIGHT.

COMMON MISCONCEPTIONS ABOUT FREE FALL

ONE OF THE MOST PERSISTENT MISCONCEPTIONS ABOUT FREE FALL IS THAT HEAVIER OBJECTS FALL FASTER THAN LIGHTER ONES. AS WE'VE DISCUSSED, THIS IS ONLY TRUE WHEN AIR RESISTANCE IS A SIGNIFICANT FACTOR. IN A VACUUM, ALL OBJECTS ACCELERATE AT THE SAME RATE DUE TO GRAVITY, REGARDLESS OF THEIR MASS. THIS WAS A REVOLUTIONARY IDEA WHEN GALILEO FIRST PROPOSED IT AND IS A TESTAMENT TO THE FACT THAT OUR EVERYDAY INTUITION CAN SOMETIMES BE MISLEADING WHEN DEALING WITH PHYSICS.

ANOTHER COMMON MISUNDERSTANDING IS THAT OBJECTS IN ORBIT ARE SOMEHOW "ABOVE" GRAVITY. THIS IS INCORRECT. OBJECTS IN ORBIT ARE VERY MUCH UNDER THE INFLUENCE OF GRAVITY; IT IS THE VERY FORCE THAT KEEPS THEM IN ORBIT. THE PERCEIVED LACK OF WEIGHT IS DUE TO THE CONTINUOUS STATE OF FREE FALL, NOT THE ABSENCE OF GRAVITY. UNDERSTANDING THESE DISTINCTIONS IS KEY TO A SOLID GRASP OF PHYSICS.

DOES FREE FALL REQUIRE DROPPING AN OBJECT?

NO, FREE FALL DOES NOT EXCLUSIVELY REQUIRE DROPPING AN OBJECT. ANY OBJECT THAT IS MOVING SOLELY UNDER THE INFLUENCE OF GRAVITY IS IN FREE FALL. THIS INCLUDES OBJECTS THROWN UPWARDS, AS THEY ARE STILL SUBJECT TO GRAVITY'S PULL ONCE THEY LEAVE THE HAND, OR OBJECTS MOVING HORIZONTALLY, AS LONG AS GRAVITY IS THE ONLY SIGNIFICANT FORCE ACTING ON THEM. THE MOTION IS DEFINED BY THE FORCES ACTING UPON IT, NOT NECESSARILY THE INITIAL ACTION THAT SET IT IN MOTION.

IS THERE GRAVITY IN SPACE?

YES, THERE IS GRAVITY IN SPACE. GRAVITY IS A UNIVERSAL FORCE THAT EXTENDS THROUGHOUT THE UNIVERSE. THE STRENGTH OF GRAVITY DECREASES WITH DISTANCE, BUT IT NEVER TRULY REACHES ZERO. ASTRONAUTS IN ORBIT AROUND EARTH EXPERIENCE GRAVITY, WHICH IS WHY THEY ARE HELD IN ORBIT. THE FEELING OF WEIGHTLESSNESS IS A RESULT OF THEIR CONTINUOUS FREE FALL, NOT AN ABSENCE OF GRAVITY. THE GRAVITATIONAL PULL OF THE EARTH IS STILL SUBSTANTIAL AT THE ALTITUDE OF THE INTERNATIONAL SPACE STATION.

FREQUENTLY ASKED QUESTIONS

Q: WHAT IS THE PRIMARY FORCE ACTING ON AN OBJECT IN FREE FALL?

A: THE PRIMARY FORCE ACTING ON AN OBJECT IN FREE FALL IS GRAVITY. ALL OTHER FORCES, SUCH AS AIR RESISTANCE, ARE ASSUMED TO BE NEGLIGIBLE IN IDEALIZED FREE FALL.

Q: DOES THE MASS OF AN OBJECT AFFECT HOW FAST IT FALLS IN FREE FALL?

A: NO, IN A VACUUM, THE MASS OF AN OBJECT DOES NOT AFFECT HOW FAST IT FALLS IN FREE FALL. ALL OBJECTS ACCELERATE AT THE SAME RATE DUE TO GRAVITY, REGARDLESS OF THEIR MASS.

Q: WHAT IS THE ACCELERATION DUE TO GRAVITY ON EARTH?

A: THE ACCELERATION DUE TO GRAVITY ON EARTH IS APPROXIMATELY 9.8 METERS PER SECOND SQUARED (m/s^2), OFTEN DENOTED BY THE SYMBOL 'G'.

Q: HOW IS FREE FALL DIFFERENT FROM PROJECTILE MOTION?

A: FREE FALL REFERS TO MOTION SOLELY UNDER GRAVITY WITH TYPICALLY NO HORIZONTAL COMPONENT OF INITIAL VELOCITY. PROJECTILE MOTION INVOLVES AN OBJECT LAUNCHED WITH AN INITIAL VELOCITY THAT HAS BOTH HORIZONTAL AND VERTICAL COMPONENTS, AND IT THEN MOVES UNDER THE INFLUENCE OF GRAVITY.

Q: CAN AN OBJECT THROWN UPWARDS BE IN FREE FALL?

A: YES, AN OBJECT THROWN UPWARDS IS IN FREE FALL ONCE IT HAS BEEN RELEASED AND IS ONLY UNDER THE INFLUENCE OF

GRAVITY. ITS MOTION WILL BE UPWARDS INITIALLY, THEN DOWNWARDS, BUT ITS ACCELERATION REMAINS CONSTANT AND DIRECTED DOWNWARDS.

Q: WHAT IS TERMINAL VELOCITY?

A: TERMINAL VELOCITY IS THE CONSTANT SPEED THAT A FREELY FALLING OBJECT EVENTUALLY REACHES WHEN THE RESISTANCE OF THE MEDIUM THROUGH WHICH IT IS FALLING PREVENTS FURTHER ACCELERATION. THIS OCCURS WHEN THE DRAG FORCE EQUALS THE FORCE OF GRAVITY.

Q: WHY DO ASTRONAUTS APPEAR TO BE WEIGHTLESS IN ORBIT?

A: ASTRONAUTS APPEAR WEIGHTLESS IN ORBIT BECAUSE THEY ARE IN A CONTINUOUS STATE OF FREE FALL AROUND THE EARTH, ALONG WITH THEIR SPACECRAFT. THEY ARE FALLING AT THE SAME RATE, SO THEY DON'T EXPERIENCE THE NORMAL SENSATION OF WEIGHT.

Q: DOES AIR RESISTANCE AFFECT FREE FALL?

A: IN IDEALIZED PHYSICS, AIR RESISTANCE IS IGNORED FOR FREE FALL. IN REALITY, AIR RESISTANCE IS A SIGNIFICANT FACTOR THAT OPPOSES THE MOTION OF FALLING OBJECTS AND CAN AFFECT THEIR SPEED AND TRAJECTORY.

Q: WHAT ARE THE KINEMATIC EQUATIONS USED TO DESCRIBE FREE FALL?

A: THE KINEMATIC EQUATIONS USED TO DESCRIBE FREE FALL ARE DERIVED FROM THE PRINCIPLES OF CONSTANT ACCELERATION AND INCLUDE FORMULAS FOR VELOCITY, DISPLACEMENT, AND TIME, WITH ACCELERATION TYPICALLY REPLACED BY 'G'.

Q: IS GRAVITY PRESENT IN SPACE?

A: YES, GRAVITY IS PRESENT IN SPACE. IT IS A UNIVERSAL FORCE THAT EXISTS BETWEEN ALL OBJECTS WITH MASS, AND IT IS RESPONSIBLE FOR PHENOMENA LIKE ORBITS AND THE MOTION OF CELESTIAL BODIES.

What Is Free Fall Motion In Physics

What Is Free Fall Motion In Physics

Related Articles

- [vibration definition physics](#)
- [uw physics building](#)
- [what is momentum and impulse in physics](#)

[Back to Home](#)